

Measure: Statin Use for Patients with Diabetes or Cardiovascular Disease

Statin Therapy

	Diabetes	Cardiovascular Disease
Population & Measurement Range	Patients 40-75 with Type I or II Diabetes dispensed at least 2 diabetic medications of any intensity during the measurement year and were dispensed a statin medication fill during the measurement year, with medication adherence of 80%	Males 21-75 and Females 40-75 during the measurement year diagnosed with atherosclerotic cardiovascular disease and dispensed at least one high or moderate intensity statin medication during the measurement year, with medication adherence of 80% OR Patients over 20 years old who have ever had a LDL-C level \geq 190 mg/dL or previous/current diagnosis of familial hypercholesterolemia
Required Exclusion Codes	Annual documentation of: Myopathy (G72); or rhabdomyolysis (M62.82); pregnancy (Z33); lactation (Z39.1); N18.6; or cirrhosis (K70.30, K74); liver disease (K76.9); polycystic ovary syndrome (E28.2) during current and prior year. Clomiphene prescription G9687	Annual documentation of: Myalgia (M79.1), myositis (M60), myopathy (G72), or rhabdomyolysis (M62.82) Pregnancy (Z33), In-vitro fertilization (O09.81), Clomiphene prescription, N18.6; or cirrhosis (K70.30, K74) during current and prior year. G2081; G2090; G2091

Captured by pharmacy claims, not included in outreach materials