

Measure:

Nephropathy

Coding:

Must document with a diagnosis code for diabetes as well as evidence of one of the four applicable codes to satisfy measure.

Nephropathy screening with a urine protein test CPT: 81000 – 81003, 81005, 82042 – 82044, 84156

Positive microalbuminuria test result documented and reviewed CPT II: 3060F or 3062F

Negative microalbuminuria test result documented and reviewed CPT II: 3061F

LOINC: 11218-5, 12842-1, 13705-9, 13801-6, 13956-5, 13992-3, 14956-7, 14957-5, 14958-3, 14959-1, 1753-3, 1754-1, 1755-8, 1757-4, 179819-4, 18373-1, 20454-5, 20621-9, 21059-1, 21482-5, 26801-1, 27298-9, 2887-8, 2888-6, 2889-4, 2890-2, 30000-4, 30001-2, 30003-8, 32209-9, 32294-1, 32551-4, 34366-5, 35663-4, 40486-3, 40662-9, 40663-7, 43605-5, 43606-3, 43607-1, 44292-1, 47558-2, 49002-9, 49023-5, 50209-6, 50561-0, 50949-7, 51190-7, 53121-0, 53525-2, 53530-2, 53531-0, 53532-8, 56553-1, 57369-1, 57735-3, 5804-0, 58448-2, 58992-9, 59159-4, 60678-0, 63474-1, 6941-9, 6942-7, 76401-9, 77253-3, 77254-1, 77940-5, 89998-9, 89999-7, 900000-1, 9318-7

Documented evidence of nephropathy with treatment CPT II: 3066F Nephropathy treatment (dialysis, ESRD, CRF, ARF, Renal Insufficiency, visit with Nephrologist),

ACE/ARB therapy CPT II: 4010F

ICD-10CM: E08.21-E08.22, E08.29, E09.21-E09.22, E09.29, E10.21-E10.22, E10.29, E11.21-E11.22, E11.29, E13.21-22, E13.29, I12.0, I12.9, I13.0, I13.10, I13.11, I13.2, I15.0, I15.1, N00.0-9, N01.0-9, N02.0-9, N03.0-9, N04.0-9, N05.0-9, N06.0-9, N07.0-9, N08, N14.0-4, N17.0-2, N17.8, N17.9, N18.1-6, N18.9, N19, N25.0, N25.1, N25.81, N25.89, N25.9, N26.1, N26.2, N26.9, Q60.0-6, Q61.00-02, Q61.11, Q61.19, Q61.2-5, Q61.8, Q61.9, R80.0-3, R80.8, R80.9

N:

Patients 18–75 with a diagnosis of Type I or Type II Diabetes.

Must document with a diagnosis code for diabetes as well as evidence of one of the four applicable codes to satisfy measure.

Excl:

Patients with documentation of ESRD, Kidney Transplant, Nephrologist visit, or Renal insufficiency. Patients with gestational diabetes.

Patients in LTC, SNF, or hospice care.

